



Impact of Intensive LDL Cholesterol Lowering on Coronary Artery Atherosclerosis Progression

A Serial CT Angiography Study

Sanghoon Shin, MD,^{a,b} Hyung-Bok Park, MD,^{b,c} Hyuk-Jae Chang, MD,^b Reza Arsanjani, MD,^d James K. Min, MD,^e Yong-Jin Kim, MD, PhD,^f Byoung Kwon Lee, MD,^g Jung-Hyun Choi, MD,^h Geu-Ru Hong, MD,^b Namsik Chung, MD^b

ABSTRACT

OBJECTIVES The aim of this study was to explore the relationship between temporal changes in coronary plaque volume and the intensity of lipid-lowering treatments, utilizing coronary computed tomography angiography (CTA).

BACKGROUND Coronary CTA has acceptable accuracy in terms of quantitative measurement of plaque volume. Although, coronary CTA is perhaps capable of identifying the differences in plaque volume progression according to the intensity of lipid lowering treatment, to date, few studies have examined this notion.

METHODS In this multicenter, observational study, the authors reviewed 467 patients who underwent serial coronary CTA with a scan period of more than 2 years (median 3.2 years [2.4 to 4.8]) apart, and whose laboratory data were available within 1 month of both the baseline and follow-up coronary CTA. Among them, 147 patients (comprising 336 vessels) with visible plaque were enrolled in this study. The authors performed quantitative assessment of coronary plaque in both. Patients who achieved a low-density lipoprotein cholesterol (LDL-C) with a cut off value below 70 mg/dl at follow-up were compared with those who did not.

RESULTS Patients with LDL-C below 70 mg/dl displayed a significant attenuation in plaque progression as compared with those with follow-up LDL-C levels ≥ 70 mg/dl (12.7 ± 38.2 mm³ vs. 44.2 ± 73.6 mm³, respectively; $p = 0.014$). In multivariate analysis, factors influencing plaque progression per year was follow-up LDL-C levels ≥ 70 mg/dl (beta 0.193; $p = 0.021$).

CONCLUSIONS Strict LDL-C control appeared to significantly attenuate plaque volume progression based on noninvasive quantitative assessment by coronary CTA. (J Am Coll Cardiol Img 2017;10:437-46)

© 2017 by the American College of Cardiology Foundation.

From the ^aDivision of Cardiology, Department of Internal Medicine, NHIS Ilsan Hospital, Goyang-si, Gyeonggi-do, South Korea; ^bDivision of Cardiology, Severance Cardiovascular Hospital, Yonsei University College of Medicine, Yonsei University Health System, Seoul, South Korea; ^cDivision of Cardiology, Department of Internal Medicine, Myungji Hospital, Goyang-si, Gyeonggi-do, South Korea; ^dMayo Clinic, Division of Cardiology, Department of Internal Medicine, Scottsdale, Arizona; ^eWeill Cornell Medical College and the New York-Presbyterian Hospital, New York, New York; ^fDivision of Cardiology, Seoul National University Hospital, Seoul, South Korea; ^gDivision of Cardiology, Gangnam Severance Hospital, Seoul, South Korea; and the ^hDivision of Cardiology, Busan Medical Center, Busan, South Korea. This work was supported by the ICT R&D program of MSIP/IITP [10044910, Development of Multi-modality Imaging and 3D Simulation-Based Integrative Diagnosis-Treatment Support Software System for Cardiovascular Diseases]. Dr. Min has received funding from the National Institutes of Health/National Heart, Lung, and Blood Institute (NIH/NIHLBI) grants R01HL111141, R01HL115150, R01HL118019, and U01HL105907, and Qatar National Research Foundation grant NPRP09-370-3-089; is a consultant for HeartFlow; is on the scientific advisory board of Arineta; has ownership of MDDX and Autoplaq; and has a research agreement with GE Healthcare. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose. Prediman K. Shah, MD, served as the Guest Editor for this paper.

Manuscript received February 19, 2016; revised manuscript received April 1, 2016, accepted April 18, 2016.

ABBREVIATIONS AND ACRONYMS

BMI = body mass index

CAD = coronary artery disease

CTA = computed tomography angiography

ECG = electrocardiogram

HDL-C = high-density lipoprotein cholesterol

IVUS = intravascular ultrasound

LDL-C = low-density lipoprotein-cholesterol

Coronary atherosclerosis is a major cause of morbidity and mortality, carrying a global burden of 17 million deaths annually (1). Low-density lipoprotein cholesterol (LDL-C) has been demonstrated to play an essential role in the progression of atherosclerotic coronary artery disease (CAD). Prior clinical trials have demonstrated a salutary effect of lowering LDL-C with statins (2,3), whereby intensive LDL-C lowering was found to be associated with improved event-free survival (4). Early studies documented that lowering LDL-C levels reduces

the progression of CAD observed via serial coronary angiography (5). Further still, the precise quantification of atheroma volume using intravascular ultrasound (IVUS) has shown that intensive lowering of LDL-C levels can halt the progression of CAD (6), or even promote atherosclerotic regression (7).

SEE PAGE 447

Serial assessment of coronary plaques using coronary angiography or IVUS has contributed to the understanding of the natural history and pathophysiology of CAD, and is therefore considered a surrogate endpoint for evaluating novel cardiovascular therapeutics (8-10). Nevertheless, these methods are invasive, expensive, and can be associated with significant adverse complications, making them unsuitable for routine clinical use. In addition, most available studies relative to invasive imaging techniques have typically included fewer numbers of patients with a generally shorter-term follow-up. Recent improvements in coronary computed tomography angiography (coronary CTA) technology have allowed for the noninvasive assessment of CAD with high diagnostic performance compared with invasive measures (11,12). In light of this, coronary CTA has emerged as a potential alternative to invasive angiography in many circumstances, including (but not confined to) stenosis grading and plaque quantification (13-15). Although a small number of serial coronary CTA studies have been published (16-18), most prior analyses were limited to a small study sample or a specific subset of lesions. Thus, in the present study, our objective was to examine changes in coronary atherosclerosis along the full length of the coronary vascular tree using coronary CTA, as well as to assess the serial changes in coronary plaque volume occurring as a result of intensive lowering of LDL-C levels.


METHODS

PATIENT SELECTION. In this multicenter, observational study, we enrolled 467 consecutive

participants who underwent serial coronary CTA using 64-slice multidetector computed tomography and whose laboratory data were available within 1 month of both the baseline and follow-up evaluations at 4 centers in Korea consisting of Severance Cardiovascular Hospital, Seoul National University Hospital, Gangnam Severance Hospital, and Busan Medical Center. Patients with plaque-free coronary arteries ($n = 153$), a history of coronary bypass surgery ($n = 108$), or inadequate imaging quality for analysis ($n = 59$) were excluded. Normal, plaque-free coronary arteries ($n = 76$), stent-implanted vessels ($n = 17$), or vessels with poor imaging quality due to artifacts ($n = 12$) were excluded from the per-vessel analysis. In total, 147 patients (comprising 336 vessels) with plaque visible using coronary CTA were enrolled in this study (Figure 1). The period between coronary CTA scans was at least 24 months. Patients were referred for coronary CTA for a variety of indications, including general health evaluation, evaluation of symptoms, and signs of cardiac disease (Table 1). The appropriate institutional review board committee approved this study.

CORONARY CTA PROTOCOL. Patients with pre-multidetector computed tomography heart rates >65 beats/min received a single oral dose (50 mg) of metoprolol tartrate (Betaloc, Yuhan, Seoul, Korea), 1 to 2 h before their CT examination unless β -adrenergic blocking agents were contraindicated (overt heart failure or atrioventricular conduction abnormalities). Patients were scanned using a 64-slice CT scanner in various centers. Coronary CTA was performed using prospective electrocardiogram (ECG) gating with the following scan parameters: rotation time 330 ms; slice collimation 64.0×0.6 mm; tube voltage 100 to 120 kV; tube current 600 to 800 mA, depending on patient size; table feed/scan 3.8 mm; and pitch factor 0.2. ECG-based tube current modulation was applied to 65% of the R-R interval. A real-time bolus-tracking technique was used to trigger scan initiation.

Contrast enhancement was achieved with 60 ml iopamidol (370 mg iodine/ml, Iopamiro, Bracco, Milan, Italy) injected at 5 ml/s, followed by an injection of 30 ml of diluted contrast (contrast agent/saline = 3:7), followed by 30 ml of saline (5 ml/s) using a power injector (Envision CT, Medrad, Indianola, Pennsylvania) via an antecubital vein. Axial images were reconstructed using retrospective ECG gating, with a slice thickness of 0.75 mm, a slice increment of 0.4 mm, and a medium-to-smooth convolution kernel. Optimal datasets with the best image quality were reconstructed, mainly during the mid-to-end



diastolic phase. The follow-up scan was performed at least 2 years after the initial CT using the same protocol.

CORONARY CTA IMAGE ANALYSIS. Datasets (baseline and follow-up) were transferred to an offline workstation for analysis using semiautomated plaque analysis software (QAngioCT Research Edition v2.0.2, Medis Medical Imaging Systems, Leiden, the Netherlands) (19). Three experienced observers (more than 3 years coronary CTA experience) blinded to the

groups analyzed all the scans and confirmed by Level III-equivalent cardiologists with experience interpreting several thousand coronary CTA scans. The major vessels (left anterior descending coronary artery, left circumflex coronary artery, and right coronary artery) were considered for analysis using the modified 17-segment American Heart Association model for coronary segment classification (20). Lesions were matched in baseline and follow-up images using branch points as landmarks. Definitions of the inner lumen and outer vessel areas were performed semiautomatically, following a stepwise approach. First, the observer placed an ostial proximal point and a distal point, and then a centerline originating from the ostium was automatically extracted. Straightened, multiplanar reformatted images were then generated, and the lumen and vessel borders were detected longitudinally by the software in 4 different cut planes and corrected by the observer. On the basis of these longitudinal contours, cross-sectional images at 0.5-mm intervals were calculated to create transverse lumen and vessel wall contours. These were

TABLE 1 Causes of Referral for Coronary CTA

	Initial	Follow-Up
Asymptomatic	83 (56.5)	93 (63.3)
Noncardiac chest pain	12 (8.2)	9 (6.1)
Atypical chest pain	42 (28.6)	33 (22.4)
Typical chest pain	5 (3.4)	8 (5.4)
Shortness of breath	5 (3.4)	4 (2.7)

Values are n (%).
 CTA = computed tomography angiography.


examined, and if necessary, adjusted by the observer. Gradient magnitude images derived from the coronary CTA images and displaying the degree of CT density change were used to verify the lumen and vessel wall borders (Figure 2, Online Figure 1).

MEASUREMENT OF CLINICAL VARIABLES AND CONFOUNDING VARIABLES. Clinical risk factors, including hypertension, diabetes mellitus, and smoking status, were systematically obtained. Height, body weight, and blood pressure were measured during the visits. Total cholesterol, triglycerides, high-density lipoprotein cholesterol (HDL-C), LDL-C, fasting plasma glucose, glycated hemoglobin, blood urea nitrogen, and serum creatinine levels were measured within 1 month of the CT scan. Diabetes was defined as treatment with oral hypoglycemic agents or insulin, or fasting glucose ≥ 126 mg/dl. Hypertension was defined as systolic blood pressure ≥ 140 mm Hg and/or diastolic blood pressure ≥ 90 mm Hg, or treatment with

antihypertensive agents. Patients were considered to be active smokers if they currently smoked or had smoked within the month before the initiation of the study. Dyslipidemia was defined as total cholesterol ≥ 240 mg/dl, LDL-C ≥ 130 mg/dl, HDL-C ≤ 40 mg/dl, triglycerides ≥ 150 mg/dl, and/or treatment with lipid-lowering agents. Laboratory values for the lipid parameters were obtained within 1 month before coronary CTA examination. The decisions to use statin therapy and the treatment target of LDL-C were determined by the individual clinician.

STATISTICAL METHODS. Continuous variables are expressed as median and interquartile range, whereas categorical variables are presented as absolute values and proportions. Differences between continuous variables were analyzed using an independent Student *t* test, whereas differences between categorical variables were analyzed by the use of a chi-square test. Changes in lesion characteristics were analyzed using a Wilcoxon signed rank test. Variables that

FIGURE 2 Representative Imaging for Lumen and Vessel Wall Contour Using Q-angio CT



Example of vessel analysis using semi-automated plaque analysis software (QAngioCT Research Edition v2.0.2, Medis Medical Imaging Systems, Leiden, the Netherlands). After the 3-dimensional centerline was generated from the computed tomography (CT) dataset, automated lumen and vessel wall contour detection was performed.

showed a significant relationship in previous studies and parameters with a p value <0.3 in the univariate analysis were defined as confounding variables related to the dependent and independent measures. The latter covariates included age, hypertension, active smoking, and follow-up LDL-C <70 mg/dl. All statistical analyses were performed using the Statistical Package for the Social Sciences version 19 (SPSS, Chicago, Illinois), and a p value <0.05 was considered significant for all analyses.

RESULTS

STUDY POPULATION. Clinical characteristics of the patients are reported in **Table 2**. The mean age was 62 ± 8 years, and 57% of the patients were male. Hypertension, diabetes mellitus, and dyslipidemia were present in 65%, 33%, and 47% of the patients, respectively. In addition, 20% of the patients were active smokers. Patients exhibiting a follow-up LDL-C <70 mg/dl had a higher prevalence of diabetes mellitus (p = 0.002). Although, there was no difference in National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) (NCEP ATP III) score and Framingham risk scores between the 2 groups, the number of risk factors were greater in the follow-up LDL <70 mg/dl group (1.95 ± 1.08 vs. 1.54 ± 0.95; p = 0.029). During a mean interscan period of 3.2 ± 1.1 years, use of statins increased considerably after baseline coronary CTA imaging (38.8% vs. 72.8%; p < 0.001), with a higher rate of use in patients achieving follow-up LDL-C <70 mg/dl (**Table 3, Figure 3**).

COMPARISON BETWEEN BASELINE AND FOLLOW-UP CHANGES IN CORONARY ARTERIES AND ATHEROSCLEROSIS PROGRESSION. Changes in coronary artery severity and atherosclerotic progression between baseline and follow-up study are displayed in **Table 4**. During this time, the lumen volume decreased (p < 0.001), whereas the vessel volume (p = 0.029), plaque volume (p < 0.001), and the proportion of dense calcified plaque (p < 0.001) increased. Furthermore, significant elevation in mean plaque burden (p < 0.001) and plaque burden at the minimal lumen area (p < 0.001) were also observed. Total plaque volume change was 36.3 ± 67.7 mm³, and the annual change in plaque volume was 13.0 ± 25.1 mm³ based on a per-patient analysis. The mean annual plaque volume progression rate was 3.63 ± 5.83% (**Figure 4**).

CORONARY ARTERY AND ATHEROSCLEROSIS CHANGES AS A FUNCTION OF LDL-C CONTROL. Patients achieving a LDL-C <70 mg/dl demonstrated a lower

TABLE 2 Baseline Patient Characteristics Split Up Based on Follow-Up LDL-C Level

	All Patients (N = 147)	F/U LDL-C <70 mg/dl (n = 37)	F/U LDL-C ≥70 mg/dl (n = 110)	p Value
Age, yrs	62 ± 8	62 ± 8	62 ± 8	0.982
Male	84 (57.1)	20 (54.1)	64 (58.2)	0.661
BMI, kg/m ²	25.1 ± 2.7	24.4 ± 2.4	25.4 ± 2.8	0.079
Hypertension	96 (65.3)	26 (70.3)	70 (63.6)	0.463
Diabetes mellitus	49 (33.3)	20 (54.1)	29 (26.4)	0.002
Active smoking	30 (20.4)	8 (21.6)	22 (21.2)	0.952
Dyslipidemia	42 (27.3)	12 (32.5)	28 (25.5)	0.409
History of PCI	7 (4.8)	3 (8.1)	4 (3.7)	0.269
History of CVA	7 (4.8)	3 (8.1)	4 (3.7)	0.269
NCEP ATP III risk	9.8 ± 7.5	8.5 ± 5.9	10.2 ± 7.9	0.232
Low (<10)	81 (55.1)	21 (56.8)	60 (54.5)	
Intermediate (10~20)	47 (32.0)	12 (32.4)	35 (31.8)	
High (>20)	19 (12.9)	4 (10.8)	15 (13.6)	
Framingham risk	10.3 ± 7.9	9.9 ± 5.8	10.5 ± 8.5	0.694
Low (<10)	92 (62.6)	22 (59.5)	70 (63.6)	
Intermediate (10~20)	42 (28.6)	13 (35.1)	29 (26.4)	
High (>20)	13 (8.8)	2 (5.4)	11 (10.0)	
Number of risk factors	1.64 ± 0.99	1.95 ± 1.08	1.54 ± 0.95	0.029


Values are mean ± SD or n (%). The mean interscan period was 3.2 ± 1.1 years.
 BMI = body mass index; CVA = cerebrovascular accident; F/U = follow-up; LDL-C = low-density lipoprotein cholesterol; NCEP ATP III = the National Cholesterol Education Program Adult Treatment Panel III; PCI = percutaneous coronary intervention.

progression in plaque volume compared with patients with follow-up LDL-C levels ≥70 mg/dl (p = 0.014). Likewise, the annual change in plaque volume was also attenuated for patients achieving low LDL-C <70 mg/dl levels (p = 0.018) (**Figure 5**). Factors associated with annual plaque progression in the univariate analysis were age, body mass index (BMI), hypertension, and follow-up LDL-C level ≥70 mg/dl.

TABLE 3 The Change of Statin Use and Laboratory Test After Index Coronary CTA

	All Patients (N = 147)	F/U LDL-C <70 mg/dl (n = 37)	F/U LDL-C ≥70 mg/dl (n = 110)	p Value
Statin use before index coronary CTA	57 (38.8)	21 (56.8)	36 (32.7)	0.009
Statin use at follow up coronary CTA	107 (72.8)	37 (100)	70 (63.6)	<0.001
Laboratory test before index coronary CTA				
T-choI, mg/dl	183 ± 35	177 ± 38	185 ± 32	0.242
HDL-C, mg/dl	48 ± 10	48 ± 9	48 ± 11	0.959
LDL-C, mg/dl	114 ± 35	106 ± 37	116 ± 32	0.125
TG, mg/dl	154 ± 8.2	150 ± 69	157 ± 88	0.689
Statin use	57 (38.8)	21 (56.8)	36 (32.7)	0.009
Laboratory test at follow up coronary CTA				
T-choI, mg/dl	162 ± 37	126 ± 15	176 ± 34	<0.001
HDL-C, mg/dl	48 ± 11	48 ± 12	47 ± 10	0.832
LDL-C, mg/dl	92 ± 32	57 ± 12	104 ± 27	<0.001
TG, mg/dl	122 ± 72	104 ± 52	126 ± 76	0.105

Values are n (%) or mean ± SD.
 HDL-C = high-density lipoprotein cholesterol; T-choI = total cholesterol; TG = triglyceride; other abbreviations as in **Tables 1 and 2**.



In per-patient multivariate analysis, follow-up LDL-C levels ≥ 70 mg/dl (beta 0.193; $p = 0.021$) were strong determinants of annual plaque progression adjusted by age, BMI, hypertension, and active smoking (Table 5). However, when the baseline plaque volume was adjusted, the plaque volume was verified as a significant factor (beta 0.324; $p < 0.001$), and the follow-up LDL ≥ 70 mg/dl was found not to be significant, but rather only showed a trend (beta 0.158; $p = 0.059$). Additionally, according to the per-vessel analysis, age, BMI, active smoking, and follow-up LDL-C ≥ 70 mg/dl were determinant factors for annual plaque progression (Online Tables 1, 2).

TABLE 4 Change in Lesion Characteristics Between Initial and Follow-Up Coronary CTA

	Initial (N = 336)	Follow Up (N = 336)	p Value
Lesion length, mm	17.0 (10.1-25.8)	17.0 (10.1-25.8)	0.486
Vessel volume, mm ³	220.8 (124.0-377.5)	223.2 (130.7-379.4)	0.616
Lumen volume, mm ³	105.9 (63.4-197.5)	97.9 (58.1-170.1)	<0.001
Plaque volume, mm ³	114.3 (63.8-200.3)	118.7 (69.1-221.6)	<0.001
Dense calcium, mm ³	10.8 (3.1-33.7)	24.4 (9.8-60.6)	<0.001
Proportion of dense calcium, %	10.0 (3.3-22.6)	22.3 (11.9-33.9)	<0.001
Mean plaque burden, %	51.7 (44.2-57.7)	55.1 (48.9-60.9)	<0.001
Maximal plaque thickness, mm	1.73 (1.37-2.12)	1.85 (1.59-2.28)	<0.001
Minimal lumen area, mm ²	5.0 (3.7-6.9)	4.5 (3.2-6.1)	<0.001
Minimal lumen diameter, mm	2.3 (1.9-2.7)	2.1 (1.7-2.5)	<0.001
Plaque burden at minimal lumen area, %	61.4 (52.4-70.5)	66.6 (59.9-74.4)	<0.001


Values are median (interquartile range).
LAD = left anterior descending coronary artery; LCX = left circumflex coronary artery; RCA = right coronary artery.

SUBGROUP ANALYSIS COMPARING STATIN USERS WHO ACHIEVED LDL-C <70 mg/dl VERSUS THOSE WHO DID NOT. A subgroup analysis on all statin users ($n = 107$), to see whether a more aggressive LDL-C approach could reduce plaque progression, was performed. All patients achieving a LDL-C <70 mg/dl took statin ($n = 37$), whereas 63% ($n = 70$) patients with follow-up LDL-C levels ≥ 70 mg/dl were on statin. Subgroup analysis of patients taking statins demonstrate that the annual plaque volume progression (4.6 ± 15.0 mm³ vs. 14.5 ± 22.0 mm³; $p = 0.015$) could be attenuated with aggressive LDL control (Table 6).

DISCUSSION

The present study demonstrated that achievement of LDL-C control to levels <70 mg/dl is associated with a reduced rate and volume of plaque progression, both on a per-patient as well as per-vessel basis. Further, we observed that per-patient annual plaque volume increased in this near-term period on the basis of presence of age and active smoking, but not by the presence of diabetes mellitus or hypertension.

Conventionally, quantitative coronary angiography or IVUS have been employed for diagnosis of coronary atherosclerosis or coronary artery stenosis. However, a potential pitfall of these procedures is their relatively expensive cost as well as their inherent invasive nature (21,22). Recent improvements in CT technology have permitted characterization of plaque morphology (23,24), quantification of plaque volume (25,26), and stenosis evaluation (12,27). There have been numerous attempts to utilize coronary CTA for measuring plaque volume and to reveal other clinical implications, such as use for estimating ischemia (28). However, these attempts have been performed manually, examining 1-mm cross-sectional slices of the coronary vascular bed, requiring considerable time and effort, and making large-scale investigations difficult. Hence, most research involving plaque volume assessments by coronary CTA have examined only a small number of patients. Recently developed semiautomated plaque quantification methods allow for the measurement of plaque volumes with only minor modifications, thereby increasing the speed of assessment and allowing more patients to be enrolled in these studies. Coronary CTA utilizing these semi-automated techniques is highly reproducible (29), with plaque volumes correlating well with plaque volumes measured by IVUS (19,30). Coronary CTA has several advantages over IVUS, given its noninvasive nature, as well as overall cost. Therefore, coronary CTA may prove useful for future quantitative research into plaque progression.



LDL-C contributes to the progression of atherosclerotic CAD. Furthermore, numerous serial IVUS studies have demonstrated that strict LDL-C control can lead to plaque retardation or regression, as well as

prior research studies incorporating CT. In the ASTEROID (A Study To Evaluate the Effect of Rosuvastatin on Intravascular Ultrasound-Derived Coronary Atheroma Burden) trial (7), patients using very




TABLE 5 Association Between Annual Plaque Volume Change and Clinical Variables

	Univariate Beta	p Value	Multivariate Beta	p Value
Age, yrs	0.130	0.118	0.149	0.077
Male	0.083	0.316		
BMI, kg/m ²	0.165	0.045	0.113	0.178
Hypertension	0.170	0.040	0.143	0.089
Diabetes mellitus	-0.039	0.639		
Active smoking	0.105	0.207	0.125	0.137
Follow-up LDL-C \geq 70 mg/dl	0.195	0.018	0.193	0.021
Statin use	-0.123	0.137		

The mean interscan period was 3.2 ± 1.1 years.
Abbreviations as in Table 1.

high-intensity statin therapy achieved an average LDL-C of 60.8 mg/dl and demonstrated significant atherosclerosis regression. Similarly, Inoue et al. (18) enrolled 32 patients, with 24 individuals receiving fluvastatin treatment. These patients underwent serial coronary CTA imaging after a median treatment interval of 12 months. The study confirmed that plaque volume, especially low-attenuation plaque volume, was reduced. However, low-dose statin (10 mg) over this short period did not mediate changes in the total plaque volume. Moreover, the study omitted patients with severe calcification, thus introducing potential selection bias that may have influenced the general applicability of these findings. Recently, another serial coronary CTA study reported that statin therapy resulted in a reduced low-attenuation plaque progression compared to that observed in non-statin users (31). The present study's findings are in line with these prior studies and extend the overall literature that demonstrates the potential utility of coronary CTA, not only for clinical evaluation of atherosclerosis progression, but also for examining volumetric differences in atherosclerotic plaques over time. To our knowledge, these data represent the longest interscan period for a serial study to date, and provide information as to the longitudinal effects of therapy on atherosclerosis.

TABLE 6 Subgroup Analysis of Patients Taking Statins

	F/U LDL-C <70 mg/dl (n = 37)	F/U LDL-C \geq 70 mg/dl (n = 70)	p Value
Initial plaque volume, mm ³	274 \pm 193	359 \pm 295	0.118
Change in plaque volume, mm ³	12.7 \pm 38.2	41.8 \pm 60.0	0.009
Annual change in plaque volume, mm ³	4.6 \pm 15.0	14.5 \pm 22.0	0.015

Values are mean \pm SD.
Abbreviations as in Table 3.

The ACC/AHA guidelines (32), published in 2013, recommended basing the decision on scoring system and that patients whose overall 10-year risk score was determined to be 7.5% or greater should start statin therapy. In these guidelines, a "treat to target" goal for LDL-C is no longer recommended. However, our subgroup analysis showed that even though the patients used statin, reducing LDL-C lower than 70 mg/dl attenuated plaque progression, which means using statin is not always effective to reduce plaque progression.

Given the strong relationship between LDL-C lowering and reduction of atherosclerosis progression, the present findings indicate that noninvasive evaluation of plaque volumetric changes can be used as a surrogate marker of cardiovascular risk. Assessment of morbidity and mortality as primary endpoints in conventional large-scale clinical trials of established and novel agents is and will remain the "gold standard" assessment of therapeutic efficacy. Nevertheless, these studies are generally hindered by a considerable financial cost and lengthy follow-up duration. Invasive imaging has therefore been employed to evaluate atherosclerosis as a surrogate marker in clinical trials that are designed to investigate plaque progression. Given the safety concerns, patients enrolled in invasive imaging studies (primarily IVUS) have been selected as a function of the disease burden, with prior investigations requiring individuals to have one obstructed vessel and another diseased coronary vessel that has not undergone angioplasty. Therefore, only patients with advanced atherosclerotic, obstructive disease have traditionally been enrolled in IVUS studies. IVUS studies remain expensive and invasive, and are inappropriate for routine serial follow-up studies. By contrast, we were able to evaluate patients with mild and moderate plaque burden in our current study, based on the noninvasive nature of coronary CTA, which permits plaque quantification with accuracy similar to IVUS (19). However, a randomized prospective comparative trial between coronary CTA and IVUS will be needed.

STUDY LIMITATIONS. This was a retrospective study, which might have potentially led to selection bias. Furthermore, only baseline and final lipid data were available, and as a consequence, longitudinal LDL-C control between coronary CTAs could not be confirmed. The fact that the proportion of patients on statin during the follow-up period was higher in the LDL <70 mg/dl group was a predictable result, indicating the individual clinician felt the risk to be high enough in the LDL <70 mg/dl group to set up a low target for control. However, the lack of significant correlation between statin administration and plaque

progression (Table 5) could be secondary to the fact that statin usage at a given time point was denoted to represent the entire follow-up period, which is a limitation of this retrospective study. When the baseline plaque volume was adjusted, the plaque volume was verified as a significant factor, and this attenuates the impact of follow-up LDL ≥ 70 mg/dl level, but rather only showed a trend. Therefore, it is correct that larger plaque volume at baseline leads to larger increase in plaque volume. However, there was no significant difference in the baseline plaque volume between the 2 groups, and lower LDL level, which is a modifiable factor, can markedly influence plaque progression in addition to the baseline plaque volume. Although we employed strict, standardized criteria for coronary CTA assessment, atherosclerosis findings may be influenced by the Hounsfield unit density, which is affected by a myriad of factors. Hence, decisions regarding plaque, lumen, or vessel volumes in this study were based on consensus among at least 2 physicians viewing the same data. Despite these limitations, this study used coronary CTA to estimate plaque volume in multiple patients and confirmed that plaque progression was slowed in patients who strictly controlled their LDL-C levels.

CONCLUSIONS

The present study demonstrated that achievement of LDL-C control to levels < 70 mg/dl are associated with a reduced rate and volume of plaque progression.

Therefore, strict LDL-C control results in a decrease in the progression of annual plaque volumes as observed by serial coronary CTA.

ADDRESS FOR CORRESPONDENCE: Dr. Hyuk-Jae Chang, Division of Cardiology, Department of Internal Medicine, Yonsei University College of Medicine, 50 Yonsei-ro, Seodaemun-gu, Seoul 120-752, South Korea. E-mail: hjchang@yuhs.ac.

PERSPECTIVES

COMPETENCY IN MEDICAL KNOWLEDGE: This study explores the relationship between temporal changes in coronary plaque volume and the intensity of lipid-lowering treatments by serial coronary CTA. Among the patients with CAD according to coronary CTA, achievement of LDL-C control to levels < 70 mg/dl decreased the progression of annual plaque volumes as observed by serial coronary CTA. Furthermore, subgroup analysis of patients taking statins also demonstrated that the annual plaque volume progression could be attenuated with aggressive LDL control.

TRANSLATIONAL OUTLOOK: This study provides significant evidence that coronary plaque volume can be decreased by strict LDL-C control and also shows the possibility of quantitating plaque progression on the basis of serial coronary CTA. Additional studies are required to confirm that the noninvasive coronary CTA may be used as a useful tool for quantitative analysis of response to medications in the future.

REFERENCES

1. Smith SC Jr., Jackson R, Pearson TA, et al. Principles for national and regional guidelines on cardiovascular disease prevention: a scientific statement from the World Heart and Stroke Forum. *Circulation* 2004;109:3112-21.
2. Pedersen TR, Kjekshus J, Berg K, et al. Randomized trial of cholesterol-lowering in 4444 patients with coronary-heart-disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;344:1383-9.
3. Shepherd J, Cobbe SM, Ford I, et al., West of Scotland Coronary Prevention Study Group. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med* 1995;333:1301-7.
4. Cannon CP, Braunwald E, McCabe CH, et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350:1495-504.
5. Blankenhorn DH, Azen SP, Krams DM, et al. Coronary angiographic changes with lovastatin therapy. The Monitored Atherosclerosis Regression Study (MARS). *Ann Intern Med* 1993;119:969-76.
6. Nissen SE, Tuzcu EM, Schoenhagen P, et al. Effect of intensive compared with moderate lipid-lowering therapy on progression of coronary atherosclerosis: a randomized controlled trial. *JAMA* 2004;291:1071-80.
7. Nissen SE, Nicholls SJ, Sipahi I, et al. Effect of very high-intensity statin therapy on regression of coronary atherosclerosis: the ASTEROID trial. *JAMA* 2006;295:1556-65.
8. Nicholls SJ, Tuzcu EM, Sipahi I, et al. Statins, high-density lipoprotein cholesterol, and regression of coronary atherosclerosis. *JAMA* 2007;297:499-508.
9. Berry C, L'Allier PL, Gregoire J, et al. Comparison of intravascular ultrasound and quantitative coronary angiography for the assessment of coronary artery disease progression. *Circulation* 2007;115:1851-7.
10. Nissen SE, Nicholls SJ, Sipahi I, et al. Effect of very high-intensity statin therapy on regression of coronary atherosclerosis: the ASTEROID trial. *J Am Med Assoc* 2006;295:1556-65.
11. Miller JM, Rochitte CE, Dewey M, et al. Diagnostic performance of coronary angiography by 64-row CT. *N Engl J Med* 2008;359:2324-36.
12. Weustink AC, Mollet NR, Neeffjes LA, et al. Diagnostic accuracy and clinical utility of noninvasive testing for coronary artery disease. *Ann Intern Med* 2010;152:630-9.
13. Rinehart S, Vazquez G, Qian Z, Murrieta L, Christian K, Voros S. Quantitative measurements of coronary arterial stenosis, plaque geometry, and composition are highly reproducible with a standardized coronary arterial computed tomographic approach in high-quality CT datasets. *J Cardiovasc Comput Tomogr* 2011;5:35-43.
14. Leber AW, Becker A, Knez A, et al. Accuracy of 64-slice computed tomography to classify and quantify plaque volumes in the proximal coronary system: a comparative study using intravascular ultrasound. *J Am Coll Cardiol* 2006;47:672-7.
15. Achenbach S, Moselewski F, Ropers D, et al. Detection of calcified and noncalcified coronary atherosclerotic plaque by contrast-enhanced, submillimeter multidetector spiral computed

- tomography: a segment-based comparison with intravascular ultrasound. *Circulation* 2004;109:14-7.
16. Burgstahler C, Reimann A, Beck T, et al. Influence of a lipid-lowering therapy on calcified and noncalcified coronary plaques monitored by multislice detector computed tomography: results of the New Age II Pilot Study. *Invest Radiol* 2007;42:189-95.
17. Hoffmann H, Frieler K, Schlattmann P, Hamm B, Dewey M. Influence of statin treatment on coronary atherosclerosis visualised using multidetector computed tomography. *Eur Radiol* 2010;20:2824-33.
18. Inoue K, Motoyama S, Sarai M, et al. Serial coronary CT angiography-verified changes in plaque characteristics as an end point: evaluation of effect of statin intervention. *J Am Coll Cardiol Img* 2010;3:691-8.
19. Boogers MJ, Broersen A, van Velzen JE, et al. Automated quantification of coronary plaque with computed tomography: comparison with intravascular ultrasound using a dedicated registration algorithm for fusion-based quantification. *Eur Heart J* 2012;33:1007-16.
20. Austen WG, Edwards JE, Frye RL, et al. A reporting system on patients evaluated for coronary artery disease. Report of the Ad Hoc Committee for Grading of Coronary Artery Disease, Council on Cardiovascular Surgery, American Heart Association. *Circulation* 1975;51:5-40.
21. De Feyter PJ, Serruys PW, Davies MJ, Richardson P, Lubsen J, Oliver MF. Quantitative coronary angiography to measure progression and regression of coronary atherosclerosis: value, limitations, and implications for clinical trials. *Circulation* 1991;84:412-23.
22. Mintz GS, Garcia-Garcia HM, Nicholls SJ, et al. Clinical expert consensus document on standards for acquisition, measurement and reporting of intravascular ultrasound regression/progression studies. *EuroIntervention* 2011;6:1123-30, 9.
23. Motoyama S, Kondo T, Sarai M, et al. Multislice computed tomographic characteristics of coronary lesions in acute coronary syndromes. *J Am Coll Cardiol* 2007;50:319-26.
24. Hoffmann U, Moselewski F, Nieman K, et al. Noninvasive assessment of plaque morphology and composition in culprit and stable lesions in acute coronary syndrome and stable lesions in stable angina by multidetector computed tomography. *J Am Coll Cardiol* 2006;47:1655-62.
25. Motoyama S, Sarai M, Harigaya H, et al. Computed tomographic angiography characteristics of atherosclerotic plaques subsequently resulting in acute coronary syndrome. *J Am Coll Cardiol* 2009;54:49-57.
26. Papadopoulou SL, Neeffjes LA, Garcia-Garcia HM, et al. Natural history of coronary atherosclerosis by multislice computed tomography. *J Am Coll Cardiol Img* 2012;5 Suppl 3:S28-37.
27. Shin S, Kim KJ, Chang HJ, et al. Impact of serum calcium and phosphate on coronary atherosclerosis detected by cardiac computed tomography. *Eur Heart J* 2012;33:2873-81.
28. Nakazato R, Shalev A, Doh J-H, et al. Aggregate plaque volume by coronary computed tomography angiography is superior and incremental to luminal narrowing for diagnosis of ischemic lesions of intermediate stenosis severity. *J Am Coll Cardiol* 2013;62:460-7.
29. Papadopoulou SL, Garcia-Garcia HM, Rossi A, et al. Reproducibility of computed tomography angiography data analysis using semiautomated plaque quantification software: implications for the design of longitudinal studies. *Int J Cardiovasc Imaging* 2013;29:1095-104.
30. Papadopoulou SL, Neeffjes LA, Schaap M, et al. Detection and quantification of coronary atherosclerotic plaque by 64-slice multidetector CT: a systematic head-to-head comparison with intravascular ultrasound. *Atherosclerosis* 2011;219:163-70.
31. Zeb I, Li D, Nasir K, et al. Effect of statin treatment on coronary plaque progression - A serial coronary CT angiography study. *Atherosclerosis* 2013;231:198-204.
32. Goff DC, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk. *J Am Coll Cardiol* 2014;63:2935-59.

KEY WORDS atherosclerosis, coronary computed tomography angiography, coronary plaque quantification, low-density lipoprotein-cholesterol

APPENDIX For a supplemental figure and tables, please see the online version of this article.