

The Association of Coronary Artery Calcium With Noncardiovascular Disease



The Multi-Ethnic Study of Atherosclerosis

Catherine E. Handy, MD, MPH,^a Chintan S. Desai, MD,^a Zeina A. Dardari, MS,^a Mouaz H. Al-Mallah, MD,^b Michael D. Miedema, MD,^c Pamela Ouyang, MD,^a Matthew J. Budoff, MD,^d Roger S. Blumenthal, MD,^a Khurram Nasir, MD,^{a,e,f,g} Michael J. Blaha, MD, MPH^a

ABSTRACT

OBJECTIVES This study sought to determine if coronary artery calcium (CAC) is associated with incident noncardiovascular disease.

BACKGROUND CAC is considered a measure of vascular aging, associated with increased risk of cardiovascular and all-cause mortality. The relationship with noncardiovascular disease is not well defined.

METHODS A total of 6,814 participants from 6 MESA (Multi-Ethnic Study of Atherosclerosis) field centers were followed for a median of 10.2 years. Modified Cox proportional hazards ratios accounting for the competing risk of fatal coronary heart disease were calculated for new diagnoses of cancer, pneumonia, chronic obstructive pulmonary disease (COPD), chronic kidney disease (CKD), deep vein thrombosis/pulmonary embolism, hip fracture, and dementia. Analyses were adjusted for age; sex; race; socioeconomic status; health insurance status; body mass index; physical activity; diet; tobacco use; number of medications used; systolic and diastolic blood pressure; total and high-density lipoprotein cholesterol; antihypertensive, aspirin, and cholesterol medication; and diabetes. The outcome was first incident noncardiovascular disease diagnosis.

RESULTS Compared with those with CAC = 0, those with CAC >400 had an increased hazard of cancer (hazard ratio [HR]: 1.53; 95% confidence interval [CI]: 1.18 to 1.99), CKD (HR: 1.70; 95% CI: 1.21 to 2.39), pneumonia (HR: 1.97; 95% CI: 1.37 to 2.82), COPD (HR: 2.71; 95% CI: 1.60 to 4.57), and hip fracture (HR: 4.29; 95% CI: 1.47 to 12.50). CAC >400 was not associated with dementia or deep vein thrombosis/pulmonary embolism. Those with CAC = 0 had decreased risk of cancer (HR: 0.76; 95% CI: 0.63 to 0.92), CKD (HR: 0.77; 95% CI: 0.60 to 0.98), COPD (HR: 0.61; 95% CI: 0.40 to 0.91), and hip fracture (HR: 0.31; 95% CI: 0.14 to 0.70) compared to those with CAC >0. CAC = 0 was not associated with less pneumonia, dementia, or deep vein thrombosis/pulmonary embolism. The results were attenuated, but remained significant, after removing participants developing interim nonfatal coronary heart disease.

CONCLUSIONS Participants with elevated CAC were at increased risk of cancer, CKD, COPD, and hip fractures. Those with CAC = 0 are less likely to develop common age-related comorbid conditions, and represent a unique population of "healthy agers." (J Am Coll Cardiol Img 2016;9:568-76) © 2016 by the American College of Cardiology Foundation.

From ^aThe Johns Hopkins Ciccarone Center for the Prevention of Heart Disease, Baltimore, Maryland; ^bHeart and Vascular Institute, Henry Ford Hospital, Detroit, Michigan; ^cMinneapolis Heart Institute and Minneapolis Heart Institute Foundation, Minneapolis, Minnesota; ^dDivision of Cardiology, Los Angeles Biomedical Research Institute at Harbor-UCLA, Torrance, California; ^eCenter for Healthcare Advancement & Outcomes, Baptist Health South Florida, Miami, Florida; ^fDepartment of Medicine, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida; and the ^gDepartment of Epidemiology, Robert Stempel College of Public Health, Florida International University, Miami, Florida. This research was supported by contracts N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, and N01-HC-95169 from the National Heart, Lung, and Blood Institute, and by grants UL1-TR-000040 and UL1-TR-001079 from the National Center for Research Resources. Dr. Nasir is on the Advisory Board of Quest Diagnostic and is a consultant for Regeneron. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose. Drs. Handy and Desai contributed equally to this work.

Manuscript received August 4, 2015; revised manuscript received August 31, 2015, accepted September 3, 2015.

Over the last 4 decades, there has been a substantial decline in cardiovascular disease (CVD) mortality, partly from improved identification and modification of risk factors (1). The rate of non-CVD deaths has not declined as rapidly (2), resulting in CVD decreasing to the second leading cause of death in certain populations (3).

Coronary artery calcium (CAC) scoring is a noninvasive, direct measure of coronary atherosclerosis and powerful predictor of incident CVD and all-cause mortality (4). CAC scores increase with chronologic age (5). However, there is considerable heterogeneity within age groups (6) translating into an “equivalent” chronologic age varying up to 30 years (7). This potentially permits CAC scores to reclassify risk independent of age (8). Indeed, CAC retains a strong predictive value for all-cause mortality beyond age (9), raising the idea that CAC scores may provide a superior estimate of “arterial age” (10).

SEE PAGE 577

CAC scores have been associated with the presence of traditional risk factors and likely partially represent the cumulative burden of risk factor exposure. However, even those with no traditional CV risk factors can have elevated CAC scores and increased risk of CVD (11). Higher levels of CAC are also associated with an increased risk of death from non-CVD causes (12). This may be from higher incidences of other age-related diseases, such as cancer, which has been associated with very high CAC scores (13). There have been mixed results when examining the relationship between CAC and other non-CVDs, such as kidney disease (14,15) and decreased bone density (16,17).

To our knowledge, there have been no studies examining the relationship between CAC and other age-related diseases associated with significant morbidity and mortality, such as osteoporotic fractures, pneumonia, dementia, deep vein thrombosis (DVT), or pulmonary embolism (PE). We sought to evaluate whether CAC, as a marker of arterial age, is an independent predictor of non-CVD diagnoses. We hypothesized that having CAC = 0 would be associated with a low risk of disease and having elevated CAC would be associated with a higher burden of non-CVD.

METHODS

STUDY POPULATION. Participants were from MESA (Multi-Ethnic Study of Atherosclerosis) (18). MESA is a prospective, observational cohort of 6,814 people from 6 U.S. cities, between 45 and 84 years old and of diverse backgrounds, who were free of CVD and not under active cancer treatment. Baseline lifestyle

characteristics and anthropometric measurements were measured at the initial examination. All participants with CAC scanning at baseline and follow-up data were eligible for inclusion. Institutional Review Board approval was obtained at each site. Each participant gave written informed consent (<http://www.mesa-nhlbi.org>).

CAC SCORE MEASUREMENTS. Detailed methods for computed tomography scan technique and interpretation were previously described (19). Chicago, Los Angeles, and New York used cardiac-gated electron-beam computed tomography scanning. The other sites used multidetector computed tomography scanning. At the baseline examination, each participant was scanned 2 consecutive times with mean Agatston score used for the analysis. All images were interpreted at a single center, with good intrareader and inter-reader agreement ($\kappa = 0.92$).

FOLLOW-UP. Follow-up occurred every 9 to 12 months for a median of 10.2 years (mean 9.5 years; interquartile range: 9.7 to 10.7 years). New diagnoses were verified by review of hospital records and death certificates. Coronary heart disease (CHD)-related endpoints were adjudicated and classified by 2 physicians from the MESA mortality and morbidity review committee. Non-CVD diagnoses were abstracted from inpatient records by International Classification of Diseases-9 codes. Codes related to the following broader groups were included: chronic kidney disease (CKD) and indicators of end-stage renal failure, any malignant neoplasm, dementia, hip fracture, DVT or PE, pneumonia, and chronic obstructive pulmonary disease (COPD). A full list of codes used is in [Online Table 1](#).

STATISTICAL ANALYSIS. We modeled CAC as a continuous and binary variable (present/absent and zero or >400). In accordance with prior reports, CAC was analyzed as a continuous variable using base-2 logarithm of coronary calcium score plus 1 ($\log_2 [CAC+1]$) to determine how the doubling of calcium score affects risk and to include those with CAC = 0 (20). There were no deviations from the linear assumption when modeling continuous CAC.

Baseline characteristics are presented by CAC stratum. Continuous variables are presented as mean \pm SD, whereas categorical variables as the number with the attribute (percentage of total). Analysis of variance was used to test means across groups for normally distributed variables and Kruskal-Wallis for not normally distributed variables. Chi-square analysis was used to test differences in distributions for categorical variables.

ABBREVIATIONS AND ACRONYMS

CAC	= coronary artery calcium
CHD	= coronary heart disease
CI	= confidence interval
CKD	= chronic kidney disease
COPD	= chronic obstructive pulmonary disease
CVD	= cardiovascular disease
DVT	= deep vein thrombosis
HR	= hazard ratio
PE	= pulmonary embolism
SES	= socioeconomic status

The risk of each non-CVD diagnosis was analyzed individually and with a composite measure of the first occurrence of any one of the diagnoses. Raw proportions experiencing each diagnosis were calculated by CAC stratum and compared using chi-square analysis. Kaplan-Meier curves for non-CVD-free survival were constructed. Hazard ratios (HRs) and 95% confidence intervals (CIs) for non-CVD and CVD-free survival were then calculated for each CAC definition with sequentially adjusted modified Cox proportional hazards regression models. To account for the competing risk of a fatal CHD event, regression models were modified, which creates informative censoring because of the strong relationship between

CAC and CHD. Other non-CVD diagnoses were not considered as competing risks.

Model 1 was unadjusted. In Model 2, analyses were adjusted for age, sex, race, health insurance status, and socioeconomic status (SES) using education and income level. Because age varied between CAC groups, additional strategies were used to limit residual confounding. For each outcome, sequential fractional polynomial terms for age were fit (using the flexible “fp” command in Stata [StataCorp, College Station, Texas]), and retained throughout all outcome-specific models if their inclusion resulted in significant improvement in model variance. An age² term was also empirically tested for each outcome; however, this did not further improve model fit beyond the best fit fractional polynomial age term. Additionally, age × CAC interaction terms were tested and retained for outcomes in which they were statistically significant.

Model 3 added adjustments for body mass index; physical activity; diet; and smoking status, including never, former, and current use and pack-years of smoking. Descriptions of physical activity and diet are included in the [Online Appendix](#). Model 4 added the total number of medications used as a surrogate for burden of comorbid illness and access to care. Model 5 added adjustments for traditional cardiovascular risk factors including systolic and diastolic blood pressure, use of antihypertensive and lipid-lowering medications, total and high-density lipoprotein cholesterol, use of aspirin, and presence of diabetes.

Model 6 added a sensitivity analysis where participants with any nonfatal CHD event at or before the time of the non-CVD diagnosis were removed from the sample ([Online Table 2](#)). This accounted for the potential identification bias of a comorbid illness during an admission for CHD. Additional sensitivity analyses were performed stratifying by age category (< and ≥ 65). Statistical analysis was done using Stata 13 (StataCorp). A p value <0.05 was considered statistically significant.

RESULTS

BASELINE CHARACTERISTICS. Baseline characteristics are presented in [Table 1](#). Among the 6,814 participants, 50.1% (n = 3,416) had CAC = 0, 39.9% (n = 2,721) had CAC scores 1 to 400, and 9.9% (n = 677) had CAC >400. The mean age was 62 years with 52.9% female, 38.5% Caucasian, 27.8% African American, 22% Hispanic, and 11.8% Chinese. Those with CAC = 0 were more likely to be younger, female, and more physically active.

TABLE 1 Baseline Characteristics of Participants by CAC Stratum

	All Participants (n = 6,814)	CAC = 0 (n = 3,416)	CAC 1-400 (n = 2,721)	CAC >400 (n = 677)	p Value
Age, yrs	62.15 ± 10.2	57.97 ± 9.1	65.32 ± 9.6	70.50 ± 7.8	<0.001
Female	3,601 (52.9)	2,167 (63.4)	1,232 (45.3)	202 (29.8)	<0.001
Race/ethnicity					<0.001
Caucasian	2,622 (38.5)	1,127 (33.0)	1,144 (42.0)	351 (51.9)	
Chinese	803 (11.8)	399 (11.7)	352 (12.9)	52 (7.7)	
African American	1,893 (27.8)	1,072 (31.4)	669 (24.6)	152 (22.5)	
Hispanic	1,496 (22.0)	818 (24.0)	556 (20.4)	122 (18.0)	
Socioeconomic status					
Completed high school	5,566 (82.0)	2,808 (82.6)	2,206 (81.3)	552 (81.7)	0.42
Income level	3,232 (49.4)	1,723 (52.3)	1,215 (46.6)	294 (45.9)	<0.001
% with no health insurance	609 (8.97)	379 (11.14)	204 (7.51)	26 (3.85)	<0.001
BMI, kg/m ²	28.0 ± 5.5	28.0 ± 5.7	28.0 ± 5.4	29.0 ± 5.0	0.49
Moderate and vigorous physical activity, MET-min/wk	5,749 ± 5,896	6,013 ± 5,822	5,596 ± 6,136	5,036 ± 5,166	<0.001
Healthy diet	2,824 (45.3)	1,340 (43.2)	1,176 (47.1)	308 (48.5)	0.004
Smoking status					<0.001
Never	3,418 (50.3)	1,905 (56.0)	1,248 (46.0)	265 (39.2)	
Former	2,487 (36.6)	1,045 (30.7)	1,113 (41.0)	329 (48.7)	
Current	887 (13.1)	451 (13.3)	354 (13.0)	82 (12.1)	
Pack-years of smoking	11.3 (20.9)	7.94 (16.5)	13.42 (22.58)	19.84 (28.64)	<0.001
Systolic blood pressure, mm Hg	127 ± 21	122 ± 21	130 ± 22	135 ± 22	<0.001
Diastolic blood pressure, mm Hg	72 ± 10	71 ± 10	72 ± 10	73 ± 10	<0.001
Total cholesterol, mg/dl	194.0 ± 35.7	194.0 ± 35.0	195.0 ± 36.2	192.0 ± 37.3	0.051
HDL-C, mg/dl	51.0 ± 14.8	53.0 ± 15.0	50.0 ± 14.4	49.0 ± 14.6	<0.001
Antihypertensive therapy	2,536 (37)	984 (29)	1,161 (43)	391 (58)	<0.001
Lipid-lowering therapy	1,100 (16)	360 (11)	543 (20)	197 (29)	<0.001
Average total number of medications	3.25 ± 2.9	2.90 ± 2.7	3.43 ± 2.9	4.26 ± 3.2	<0.001
Presence of diabetes	859 (12.7)	318 (9.4)	387 (14.3)	154 (22.9)	<0.001
Aspirin use	1,298 (19.9)	462 (14.2)	607 (23.1)	229 (35.5)	<0.001

Values are mean ± SD or n (%).

BMI = body mass index; CAC = coronary artery calcium; HDL-C = high-density lipoprotein cholesterol.

CANCER. There were 710 new cancer diagnoses. The proportions of participants with cancer by CAC stratum were 6.67%, 12.6%, and 20.4% for CAC = 0, CAC 1 to 400, and CAC >400, respectively ($p < 0.001$) (Figure 1). Those with CAC >0 accounted for 67.9% of cancer cases. Unadjusted and adjusted HRs are in Tables 2 and 3. In multivariable-adjusted models (Model 5), a doubling of CAC score was associated with a 4% increased risk of cancer (HR: 1.04; 95% CI: 1.02 to 1.07) (Table 2). Those with CAC >400 had a 53% greater hazard of cancer compared with those with CAC = 0 (HR: 1.53; 95% CI: 1.18 to 1.99) (Table 3). Those with CAC = 0 had a lower hazard of being diagnosed when compared with those with CAC >0 (HR: 0.76; 95% CI: 0.63 to 0.92) (Table 4). Prostate cancer was the most common cancer diagnosed (21% of cancer cases), followed by lung cancer (14% of cancer cases), gastrointestinal/colon cancer (13% of cancer cases), breast cancer (10% of cancer cases), skin cancer (9% of cancer cases), hematologic malignancies (6% of cancer cases), and uterine/ovarian

cancer (5% of cancer cases). After accounting for sex differences, there was no strong association between CAC score stratum and overall cancer type.

CHRONIC KIDNEY DISEASE. CKD was diagnosed in 395 participants. The proportion increased with increasing CAC stratum from 3.2% in participants with CAC = 0.0 to 13.3% in participants with CAC >400 ($p < 0.001$) (Figure 1). With adjustment for CVD risk factors, a doubling of CAC score remained significantly associated with an increased hazard of being diagnosed (HR: 1.07; 95% CI: 1.03 to 1.10) (Table 2). Those with CAC >400 were 70% more likely to develop CKD compared with those with CAC = 0 (HR: 1.70; 95% CI: 1.21 to 2.39) (Table 3) and those with CAC = 0 had a lower hazard of being diagnosed with CKD (HR: 0.77; 95% CI: 0.60 to 0.98) (Table 4).

PNEUMONIA. There were 315 cases of pneumonia. Proportions ranged from 2.8% when CAC = 0 to 10.6% in those with CAC >400 ($p < 0.001$) (Figure 1). In multivariable-adjusted models (Model 5), there




TABLE 2 Hazard of New Non-CVD and CVD Diagnosis With Doubling of CAC Score

	Log₂[CAC + 1]				
	Model 1	Model 2	Model 3	Model 4	Model 5
Cancer	1.13 (1.11-1.15)	1.05 (1.02-1.07)	1.04 (1.01-1.06)	1.03 (1.01-1.06)	1.04 (1.02-1.07)
CKD	1.18 (1.15-1.20)	1.08 (1.05-1.12)	1.08 (1.04-1.11)	1.07 (1.04-1.10)	1.07 (1.03-1.10)
Pneumonia	1.17 (1.13-1.20)	1.08 (1.04-1.12)	1.07 (1.03-1.10)	1.06 (1.02-1.09)	1.07 (1.03-1.11)
DVT/PE	1.10 (1.06-1.13)	<i>1.04 (0.99-1.08)</i>	<i>1.01 (0.97-1.06)</i>	<i>1.01 (0.96-1.05)</i>	<i>1.03 (0.98-1.08)</i>
COPD	1.19 (1.14-1.24)	1.11 (1.05-1.16)	1.09 (1.04-1.15)	1.08 (1.03-1.14)	1.10 (1.05-1.16)
Dementia	1.23 (1.18-1.28)	1.05 (1.00-1.10)	1.06 (1.003-1.11)	1.06 (1.002-1.11)	1.06 (1.01-1.12)
Hip fracture	1.22 (1.14-1.29)	1.09 (1.003-1.18)	1.11 (1.02-1.20)	1.10 (1.01-1.20)	1.10 (1.01-1.21)
Any non-CVD	1.15 (1.14-1.17)	1.06 (1.04-1.08)	1.06 (1.04-1.08)	1.05 (1.03-1.07)	1.06 (1.04-1.08)
CVD	1.26 (1.23-1.28)	1.20 (1.17-1.23)	1.19 (1.16-1.22)	1.19 (1.16-1.22)	1.17 (1.14-1.20)

Statistically significant values are in **bold** for Model 5. Nonsignificant values are in *italics*. Cox proportional hazards ratios and 95% confidence intervals for non-CVD diagnoses comparing doubling of CAC score. Model 1: unadjusted. Model 2: age (best fit model, see Methods section), sex, race, health insurance status, and socioeconomic status (completed high school and income level). Model 3: Model 2 + body mass index, physical activity, diet, smoking status, and pack-years. Model 4: Model 3 + total medications used. Model 5: Model 4 + systolic blood pressure, diastolic blood pressure, antihypertensive medications, total cholesterol, high-density lipoprotein cholesterol, lipid-lowering medications, diabetes, and aspirin use.

CKD = chronic kidney disease; COPD = chronic obstructive pulmonary disease; CVD = cardiovascular disease; DVT = deep vein thrombosis; PE = pulmonary embolism; other abbreviations as in [Table 1](#).

was a 7% increased hazard of being diagnosed with pneumonia with doubling of the CAC score (HR: 1.07; 95% CI: 1.03 to 1.11) ([Table 2](#)). Those with CAC >400 had 2 times the hazard of developing pneumonia compared with those with CAC = 0 (HR: 1.97; 95% CI: 1.37 to 2.82) ([Table 3](#)). Those with CAC = 0 did not have a significantly decreased risk of pneumonia (HR: 0.77; 95% CI: 0.58 to 1.01) when compared with those with CAC >0 ([Table 4](#)). This was associated in unadjusted models but lost after controlling for demographic and lifestyle factors (Model 3, [Table 4](#)).

DVT OR PE. DVT or PE was documented in 205 participants. Among those with CAC = 0, 1.1% were

diagnosed, 1.9% in those with CAC = 1 to 400 and 2.1% in those with CAC >400 ($p < 0.05$) ([Figure 1](#)). In unadjusted models, doubling of CAC score and CAC >400 were associated with increased hazard of DVT or PE (1.10, 1.06 to 1.13; and 2.24, 1.50 to 3.35, respectively) ([Tables 2 and 3](#), respectively). Having CAC = 0 was protective in unadjusted models (0.57, 0.44 to 0.75) ([Table 4](#)). These relationships were lost after controlling for age, sex, race, SES, and health insurance status ([Tables 2 to 4](#)). In multivariable adjusted models (Model 5), there was no association with a new diagnosis of DVT/PE and doubling of the CAC score (HR: 1.03; 95% CI: 0.98 to 1.08) ([Table 2](#)), having CAC >400 (HR: 1.07; 95% CI: 0.64 to 1.78)

TABLE 3 Hazard of New Non-CVD and CVD Diagnosis With CAC >400 Compared With Those With CAC = 0

	CAC >400 Versus CAC = 0				
	Model 1	Model 2	Model 3	Model 4	Model 5
Cancer	3.60 (2.95-4.41)	1.64 (1.29-2.07)	1.49 (1.16-1.92)	1.45 (1.13-1.86)	1.53 (1.18-1.99)
CKD	4.57 (3.55-5.89)	2.00 (1.47-2.74)	1.93 (1.39-2.70)	1.75 (1.27-2.43)	1.70 (1.21-2.39)
Pneumonia	4.72 (3.53-6.30)	2.16 (1.54-3.02)	1.88 (1.33-2.68)	1.71 (1.21-2.41)	1.97 (1.37-2.82)
DVT/PE	2.24 (1.50-3.35)	<i>1.26 (0.80-2.00)</i>	<i>0.89 (0.54-1.47)</i>	<i>0.86 (0.52-1.41)</i>	<i>1.07 (0.64-1.78)</i>
COPD	5.50 (3.64-8.32)	2.69 (1.61-4.50)	2.45 (1.44-4.18)	2.17 (1.30-3.64)	2.71 (1.60-4.57)
Dementia	6.45 (4.14-10.05)	<i>1.33 (0.81-2.19)</i>	<i>1.47 (0.88-2.47)</i>	<i>1.48 (0.87-2.50)</i>	<i>1.61 (0.93-2.79)</i>
Hip fracture	8.74 (3.93-19.45)	3.18 (1.19-8.50)	4.16 (1.48-11.73)	3.91 (1.39-10.97)	4.29 (1.47-12.50)
Any non-CVD	4.16 (3.58-4.84)	1.88 (1.57-2.24)	1.78 (1.47-2.15)	1.69 (1.40-2.04)	1.80 (1.48-2.18)
CVD	9.52 (7.65-11.86)	6.01 (4.65-7.77)	5.44 (4.16-7.10)	5.11 (3.91-6.67)	4.45 (3.36-5.88)

Statistically significant values are in **bold** for Model 5. Nonsignificant values are in *italics*. Cox proportional hazards ratios and 95% confidence intervals for non-CVD diagnoses comparing those with CAC >400 with those with CAC = 0. Model 1: unadjusted. Model 2: age (best fit model, see Methods section), sex, race, health insurance status, and socioeconomic status (completed high school and income level). Model 3: Model 2 + body mass index, physical activity, diet, smoking status, and pack-years. Model 4: Model 3 + total medications used. Model 5: Model 4 + systolic blood pressure, diastolic blood pressure, antihypertensive medications, total cholesterol, high-density lipoprotein cholesterol, lipid-lowering medications, diabetes, and aspirin use.

Abbreviations as in [Tables 1 and 2](#).

TABLE 4 Hazard of New Non-CVD Diagnosis for Those With CAC = 0 Compared With Those With CAC >0

	CAC = 0 Versus CAC >0				
	Model 1	Model 2	Model 3	Model 4	Model 5
Cancer	0.44 (0.38-0.51)	0.75 (0.64-0.89)	0.77 (0.64-0.92)	0.79 (0.66-0.94)	0.76 (0.63-0.92)
CKD	0.38 (0.31-0.46)	0.69 (0.55-0.86)	0.71 (0.56-0.90)	0.74 (0.58-0.94)	0.77 (0.60-0.98)
Pneumonia	0.40 (0.32-0.50)	0.71 (0.55-0.92)	0.78 (0.60-1.02)	0.82 (0.62-1.06)	0.77 (0.58-1.01)
DVT/PE	0.57 (0.44-0.75)	0.85 (0.62-1.17)	1.01 (0.73-1.41)	1.04 (0.75-1.45)	0.94 (0.66-1.33)
COPD	0.32 (0.23-0.45)	0.56 (0.38-0.81)	0.64 (0.43-0.96)	0.69 (0.46-1.03)	0.61 (0.40-0.91)
Dementia	0.26 (0.18-0.38)	0.87 (0.59 to1.30)	0.81 (0.54-1.21)	0.80 (0.53-1.21)	0.76 (0.49-1.17)
Hip fracture	0.17 (0.09-0.33)	0.37 (0.17-0.78)	0.30 (0.14-0.67)	0.31 (0.14-0.69)	0.31 (0.14-0.70)
Any non-CVD	0.41 (0.36-0.45)	0.71 (0.63-0.81)	0.74 (0.65-0.85)	0.76 (0.66-0.87)	0.75 (0.65-0.86)
CVD	0.21 (0.17-0.25)	0.31 (0.26-0.39)	0.34 (0.27-0.42)	0.35 (0.28-0.43)	0.38 (0.30-0.47)

Statistically significant values are in **bold** for model 5. Nonsignificant values are in *italics*. Cox proportional hazards ratios and 95% confidence intervals for non-CVD diagnoses comparing those with CAC = 0 to those with CAC >0. Model 1: unadjusted. Model 2: age (best fit model, see the Methods section), sex, race, health insurance status, and socioeconomic status (completed high school and income level). Model 3: Model 2 + body mass index, physical activity, diet, smoking status, and pack-years. Model 4: Model 3 + total medications used. Model 5: Model 4 + systolic blood pressure, diastolic blood pressure, antihypertensive medications, total cholesterol, high-density lipoprotein cholesterol, lipid-lowering medications, diabetes, and aspirin use.

Abbreviations as in [Tables 1 and 2](#).

([Table 3](#)), or having CAC = 0 (HR: 0.94; 95% CI: 0.66 to 1.33) ([Table 4](#)).

CHRONIC OBSTRUCTIVE PULMONARY DISEASE.

During follow-up, 161 participants were diagnosed with COPD. Proportions increased with increasing CAC score stratum from 1.2% in those with CAC = 0 to 5.6% in those with CAC >400 (p < 0.001) ([Figure 1](#)). In Model 5, doubling of CAC score was associated with a 10% increased hazard of COPD (HR: 1.10; 95% CI: 1.05 to 1.16) ([Table 2](#)). Those with CAC >400 were 2.7 times more likely than those with CAC = 0 (HR: 2.71; 95% CI: 1.60 to 4.57) to have a new diagnosis of COPD ([Table 3](#)). Those with CAC = 0 were approximately half as likely to be diagnosed compared with those with any CAC (HR: 0.61; 95% CI: 0.40 to 0.91) ([Table 4](#)).

DEMENTIA. Dementia was diagnosed in 119 participants and had increasing proportions with higher CAC stratum from 0.61% to 4.43% (p < 0.001) ([Figure 1](#)). After multivariable adjustment, doubling of the CAC score was associated with a 6% increased risk (HR: 1.06; 95% CI: 1.01 to 1.12) ([Table 2](#)). Comparing those with CAC >400 with those with CAC = 0, unadjusted models showed an increased HR of 6.45 (95% CI: 4.14 to 10.05) ([Table 3](#)). This relationship was lost when controlling for age, sex, race, SES, and health insurance status (Model 2, [Table 3](#)). Similarly, those with CAC = 0 compared with those with CAC >0 had a significant association in unadjusted models (HR: 0.26; 95% CI: 0.18 to 0.38) but this was lost when controlling for age, sex, race, SES, and health insurance status (Model 2, [Table 4](#)).

HIP FRACTURES. Hip fractures were the least common with 59 cases. Proportions diagnosed by CAC


category ranged from 0.26% when CAC = 0 to 1.77% in those with CAC >400 (p < 0.001) ([Figure 1](#)). In multivariable adjusted models, doubling of CAC was associated with a 10% increased risk of events (1.01 to 1.21) ([Table 2](#)). Participants with CAC >400 had a higher risk of hip fracture in multivariable adjusted modeling (Model 5) (HR: 4.29; 95% CI: 1.47 to 12.5) ([Table 3](#)). Participants with CAC = 0 had a 69% lower risk of hip fracture in multivariable adjusted modeling (HR: 0.31; 95% CI: 0.14 to 0.70) ([Table 4](#)).

AGGREGATE OUTCOME OF FIRST ANY NON-CVD DIAGNOSIS.

There were 1,238 first non-CVD diagnoses. The proportion of the first occurrence of any one of these was 11.0% for CAC = 0, 22.5% for CAC = 1 to 400, and 36.9% for CAC >400 (p < 0.001) ([Figure 2](#)). In multivariable adjusted models, doubling of the CAC score was associated with a 6% increased risk of any non-CVD diagnosis (HR: 1.06; 95% CI: 1.04 to 1.08) ([Table 2](#)). Those with CAC score >400 were 1.80 times more likely to be diagnosed with any non-CVD compared with those with CAC = 0 (HR: 1.80; 95% CI: 1.48 to 2.18) ([Table 3](#)). Participants with CAC = 0 had a lower rate of non-CVD compared with those with CAC >0 (HR: 0.75; 95% CI: 0.65 to 0.86) ([Table 4](#)). Kaplan-Meier curves for the composite and individual outcomes are shown in [Online Figure 1](#).

SENSITIVITY ANALYSIS.

The results of the sensitivity analysis are in [Online Table 2](#). Results were attenuated, with similar overall conclusions after removing participants with an interim or concurrent nonfatal CHD event at the time of diagnosis of the non-CV disease. New diagnoses of cancer, CKD, COPD, and any-non CVD event remained associated with doubling of CAC score ([Online Table 2](#)).



AGE STRATIFICATION. Results of age stratified analysis are presented in Table 5. In participants ≥ 65 , doubling of the CAC score was associated with an increased hazard of cancer (HR: 1.04; 95% CI: 1.01

TABLE 5 Age-Stratified Analysis of the Doubling of CAC Score

	Log2[CAC + 1]	
	Age <65 Yrs	Age ≥ 65 Yrs
Cancer	1.04 (1.00-1.09)	1.04 (1.01-1.07)
CKD	1.05 (0.98-1.12)	1.07 (1.03-1.11)
Pneumonia	1.04 (0.97-1.11)	1.08 (1.03-1.13)
DVT/PE	1.03 (0.95-1.11)	1.02 (0.96-1.08)
COPD	1.12 (1.00-1.25)	1.09 (1.03-1.16)
Dementia	—	1.06 (1.00-1.12)
Hip fracture	1.15 (0.94-1.41)	1.09 (0.99-1.20)
Any non-CVD event	1.04 (1.00-1.07)	1.06 (1.04-1.08)
CVD	1.19 (1.14-1.24)	1.16 (1.12-1.20)

Statistically significant values are in **bold**. Nonsignificant values are in *italics*. Cox proportional hazards ratios (and 95% confidence intervals) for those <65 and those ≥ 65 years of age. Model is adjusted for age (best fit, see the Methods section), sex, race, health insurance status, socioeconomic status (completed high school and income level), body mass index, physical activity, diet, smoking status and pack-years, total number of medications used, systolic blood pressure, diastolic blood pressure, antihypertensive medications, total cholesterol, high-density lipoprotein cholesterol, lipid-lowering medications, diabetes, and aspirin use. The interaction of CAC with age < or ≥ 65 years was not significant. Abbreviations as in Tables 1 and 2.

to 1.07), CKD (HR: 1.07; 95% CI: 1.03 to 1.11), pneumonia (HR: 1.08; 95% CI: 1.03 to 1.13), COPD (HR: 1.09; 95% CI: 1.03 to 1.16), and the composite outcome (HR: 1.06; 95% CI: 1.04 to 1.08). Limited power prevented detailed analysis of modeling in those <65, although point estimates were similar with nonsignificant interaction with the age < or ≥ 65 term.

DISCUSSION

Our results demonstrate that CAC scores are associated with non-CVD diagnoses, notably cancer, CKD, COPD, and hip fractures. Participants with CAC scores >400 had a 53% adjusted increased hazard of cancer diagnosis, 70% increased hazard of CKD diagnosis, 197% increased hazard of developing pneumonia, 271% increased hazard of COPD diagnosis, and 429% increased hazard of hip fracture. Participants with CAC = 0 (50% of the population) seemed at reduced risk of non-CVD, independent of known CVD risk factors. Although these results were attenuated when excluding patients with interim nonfatal CHD events, overall conclusions remained unchanged. Taken together, higher CAC scores were highly associated with the initial development of any first non-CVD diagnosis. From a public health perspective, it is notable that 20% of the first occurrences of a non-CVD event occurred in the 10% of patients with CAC >400 and 70% of the events were in those with CAC >0.

Strengths of our analysis include the large sample, multiethnic makeup, detailed baseline risk factor assessment, and the use of a competing risk model, which is important given the known association of CAC with fatal CHD. In addition, rather than assuming a linear association, we allowed flexible modeling of age to achieve the best fit and thus limit residual confounding from age.

It is important to consider the possible mechanisms underlying these results. CAC per se is unlikely to be causally associated with non-CVD. In general, CAC is considered to be a risk integrator, and may reflect lifetime exposure to both measured and unmeasured risk factors shared in common between CVD and non-CVD outcomes (21). It may also represent a common underlying vulnerability to the development of tissue injury at sites with varying relationships to the cardiovascular system. As such, the CAC score may be one marker of an individual's risk of future disease burden.

Pneumonia was not associated with CAC scores in models removing individuals with an interim nonfatal CHD event, suggesting it may be a more

acute event that is more likely to be an incidental codiagnosis during the inpatient care for, or related to complications of, a CHD event. Furthermore, DVT/PE were weakly associated with CAC and are well known to have a variety of more acute-onset risk factors, including trauma and surgery (22), which could be less related to CVD and arterial aging. Dementia likely had too few cases to confirm an association with CAC given the large CI.

IMPLICATIONS. We hypothesized that CAC scores could be a surrogate measure of arterial aging and identify those at increased risk of age-related, non-CV disease. CAC has been described as a measure of so-called “biologic age” (7), supported by studies showing absolute CAC scores as better predictors of CVD events than age alone (10). Our results support CAC as a marker for other age-related diseases and support limited prior findings showing an association between elevated CAC scores and non-CV diseases including cancer (13) and COPD (23). Although CAC has been shown to be associated with all-cause mortality (24-26), this association may be driven by both cardiovascular and select noncardiovascular causes. The risk ratio of cardiovascular versus non-CVD diagnoses likely varies as a function of the CAC score and future mortality studies may benefit from ascertainment of cause of death and modeling the risk of both cardiovascular and noncardiovascular causes.

An important finding is that people with CAC = 0 seem to be protected from CVD and other chronic diseases. These “healthy agers” are at very low risk of CVD and non-CVD morbidity and mortality (24,27,28). There are broad implications to identifying and studying such people in the population. These people are extremely unlikely to be high users of healthcare services because of the decreased risk of comorbidities. In contrast, those with elevated CAC are at high risk of developing multiple chronic conditions and are likely to require increased services. These findings, which mirror the observation that individuals with multiple chronic conditions use a disproportionate share of health care resources (29), have marked implications for patient-centered medical homes, accountable care organizations, insurers, public health officials, and other stakeholders in the distribution of health care resource expenditures.

STUDY LIMITATIONS. New non-CVD diagnoses in MESA were obtained via International Classification of Disease hospital coding and therefore must be interpreted as time-to-diagnosis, not necessarily time-to-disease onset. This would mostly affect the

diagnoses of cancer, COPD, and CKD, which could be detected before hospitalization. Furthermore, by using inpatient codes, there may be bias toward capturing severe cases and missing milder cases that are exclusively managed in the outpatient setting. Additionally, outcome ascertainment was administrative and does not represent systematically adjudicated events as is the case for CVD-related outcomes in MESA.

The numbers of participants with hip fractures, DVT/PE, and dementia were small and our study may not be powered to firmly establish the relationship. Further studies of these outcomes are needed, perhaps leveraging electronic medical record and claims files databases that would allow examination of a larger sample of patients. In addition, follow-up time in MESA may not be sufficient for detecting risk in participants with the lowest CAC scores. At this time, our data are not powered for stratifying results based on sex or race.

CONCLUSIONS

Elevated CAC scores are associated with age-related diseases including cancer, COPD, CKD, and hip fractures. Those with CAC = 0 are less likely to develop common age-related comorbid conditions, and represent a unique population of “healthy agers.”

ACKNOWLEDGMENTS The authors thank the other investigators, the staff, and the participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://www.mesa-nhlbi.org>.

REPRINT REQUESTS AND CORRESPONDENCE: Dr. Michael J. Blaha, The Johns Hopkins Ciccarone Center for the Prevention of Heart Disease, Carnegie 565A, 600 North Wolfe Street, Baltimore, Maryland 21287. E-mail: mblaha1@jhmi.edu.

PERSPECTIVES

COMPETENCY IN MEDICAL KNOWLEDGE: CAC scoring is a noninvasive, direct measure of coronary atherosclerosis.

COMPETENCY IN PATIENT CARE AND PROCEDURAL SKILLS: CAC retains a strong predictive value for cardiovascular and noncardiovascular disease events beyond age.

TRANSLATIONAL OUTLOOK: People with CAC = 0 seem to be protected from multiple chronic noncardiovascular diseases.

REFERENCES

1. Ford ES, Ajani UA, Croft JB, et al. Explaining the decrease in U.S. deaths from coronary disease, 1980-2000. *N Engl J Med* 2007;356:2388-98.
2. Araujo F, Gouvêas C, Fontes F, La Vecchia C, Azevedo A, Lunet N. Trends in cardiovascular diseases and cancer mortality in 45 countries from five continents (1980-2010). *Eur J Prev Cardiol* 2014;21:1004-17.
3. Siegel R, Naishadham D, Jemal A. Cancer statistics for Hispanics/Latinos, 2012. *CA Cancer J Clin* 2012;62:283-98.
4. Shaw LJ, Raggi P, Schisterman E, Berman DS, Callister TQ. Prognostic value of cardiac risk factors and coronary artery calcium screening for all-cause mortality. *Radiology* 2003;228:826-33.
5. McClelland RL, Chung H, Detrano R, Post W, Kronmal RA. Distribution of coronary artery calcium by race, gender, and age: Results from the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation* 2006;113:30-7.
6. Tota-Maharaj R, Blaha MJ, McEvoy JW, et al. Coronary artery calcium for the prediction of mortality in young adults 75 years old. *Eur Heart J* 2012;33:2955-62.
7. Shaw LJ, Raggi P, Berman DS, Callister TQ. Coronary artery calcium as a measure of biologic age. *Atherosclerosis* 2006;188:112-9.
8. Grundy SM. Coronary plaque as a replacement for age as a risk factor in global risk assessment. *Am J Cardiol* 2001;88:8E-11E.
9. Tota-Maharaj R, Blaha MJ, Blankstein R, et al. Association of coronary artery calcium and coronary heart disease events in young and elderly participants in the multi-ethnic study of atherosclerosis: a secondary analysis of a prospective, population-based cohort. *Mayo Clin Proc* 2014;89:1350-9.
10. McClelland RL, Nasir K, Budoff M, Blumenthal RS, Kronmal RA. Arterial age as a function of coronary artery calcium (from the Multi-Ethnic Study of Atherosclerosis [MESA]). *Am J Cardiol* 2009;103:59-63.
11. LaMonte MJ, FitzGerald SJ, Church TS, et al. Coronary artery calcium score and coronary heart disease events in a large cohort of asymptomatic men and women. *Am J Epidemiol* 2005;162:421-9.
12. Desai CS, Ning H, Kang J, et al. Competing cardiovascular outcomes associated with subclinical atherosclerosis (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol* 2013;111:1541-6.
13. Chen WT, Huang JH, Hsieh MH, Chen YJ. Extremely high coronary artery calcium score is associated with a high cancer incidence. *Int J Cardiol* 2012;155:474-5.
14. Fox CS, Larson MG, Keyes MJ, et al. Kidney function is inversely associated with coronary artery calcification in men and women free of cardiovascular disease: the Framingham Heart Study. *Kidney Int* 2004;66:2017-21.
15. Parikh NI, Hwang SJ, Larson MG, et al. Indexes of kidney function and coronary artery and abdominal aortic calcium (from the Framingham Offspring study). *Am J Cardiol* 2008;102:440-3.
16. Bakhireva LN, Barrett-Connor EL, Laughlin GA, Kritzer-Silverstein D. Differences in association of bone mineral density with coronary artery calcification in men and women: the Rancho Bernardo study. *Menopause* 2005;12:691-8.
17. Hyder JA, Allison MA, Wong N, et al. Association of coronary artery and aortic calcium with lumbar bone density: the MESA abdominal aortic calcium study. *Am J Epidemiol* 2009;169:186-94.
18. Bild DE, Bluemke DA, Burke GL, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol* 2002;156:871-81.
19. Carr JJ, Nelson JC, Wong ND, et al. Calcified coronary artery plaque measurement with cardiac CT in population-based studies: standardized protocol of Multi-Ethnic Study of Atherosclerosis (MESA) and coronary artery risk development in young adults (CARDIA) study. *Radiology* 2005;234:35-43.
20. Detrano R, Guerci AD, Carr JJ, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;358:1336-45.
21. Blaha MJ, Silverman MG, Budoff MJ. Is there a role for coronary artery calcium scoring for management of asymptomatic patients at risk for coronary artery disease?: Clinical risk scores are not sufficient to define primary prevention treatment strategies among asymptomatic patients. *Circ Cardiovasc Imaging* 2014;7:398-408; discussion 408.
22. Anderson FA, Jr., Spencer FA. Risk factors for venous thromboembolism. *Circulation* 2003;107:19-16.
23. Park HY, Lim SY, Hwang JH, et al. Lung function, coronary artery calcification, and metabolic syndrome in 4905 Korean males. *Respir Med* 2010;104:1326-35.
24. Budoff MJ, Shaw LJ, Liu ST, et al. Long-term prognosis associated with coronary calcification: observations from a registry of 25,253 patients. *J Am Coll Cardiol* 2007;49:1860-70.
25. Shaw LJ, Giambone AE, Blaha MJ, et al. Long-term prognosis after coronary artery calcification testing in asymptomatic patients: a cohort study. *Ann Intern Med* 2015;163:14-21.
26. Raggi P, Gongora MC, Gopal A, Callister TQ, Budoff M, Shaw LJ. Coronary artery calcium to predict all-cause mortality in elderly men and women. *J Am Coll Cardiol* 2008;52:17-23.
27. Blaha M, Budoff MJ, Shaw LJ, et al. Absence of coronary artery calcification and all-cause mortality. *J Am Coll Cardiol Img* 2009;2:692-700.
28. Budoff MJ, McClelland RL, Nasir K, et al. Cardiovascular events with absent or minimal coronary calcification: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am Heart J* 2009;158:554-61.
29. Vogeli C, Shields AE, Lee TA, et al. Multiple chronic conditions: prevalence, health consequences, and implications for quality, care management, and costs. *J Gen Intern Med* 2007;22 Suppl 3:391-5.

KEY WORDS aging, biologic aging, cancer, coronary artery calcium, coronary artery disease

APPENDIX For supplemental tables and figures, please see the online version of this article.